



Aftercare Liquid Gold

During the first 24–48 hours after treatment you will experience redness, swelling, bruising and tenderness in the treated areas. These responses are normal and resolve quickly and spontaneously over the next 2-4 days. You can apply light water-based or mineral makeup where necessary.

Drink plenty of cold water for 48 hours after the procedure. Bruising at the site of injection is also common and clears completely in 1-10 days. Please read your aftercare instructions in the aftercare pack provided - there are several things you can do to help bruising. Swelling and bruising can give the appearance of 'unevenness' in the area treated. This will settle once the swelling and bruising has gone. Please contact the clinic for advice if there is any 'unevenness' present after 14 days.

After your Liquid Gold treatment you should:

- Avoid alcohol, caffeine, hot drinks and spicy food for 24 hours, as these may exacerbate bruising and swelling.
- Avoid medications such as aspirin or Nurofen for 1-2 weeks as they are blood thinning agents and may exacerbate bruising (if these medications have been advised by your GP please check with your GP before stopping them)
- Avoid supplements such as multivitamins, fish oils, glucosamine and Vitamin B or Vitamin E, which are blood thinning agents and may exacerbate bruising for 1-2 weeks (if these supplements have been advised by your GP please check with your GP before stopping them)

- Avoid exercise/exertion for 24-48 hours • Avoid excessive heat (e.g. sun exposure, hot showers, saunas, laser procedures or solarium use) for 24-48 hours
- Apply ice frequently as required to the area injected for up to 6 hours after the treatment. This will prevent or help to alleviate any bruising. It will also help to reduce any swelling. You can use Arnica cream and arnica tablets to help alleviate any bruising. Note: Bruises can last 1-10 days.
- They are TEMPORARY and will completely disappear over time with no effect on your treatment. Once bruising has formed (within the first 48 hours) it is better to apply warmth to the area to encourage the body's natural reabsorption of the bruise from the skin.
- Please advise your cosmetic practitioner prior to treatment if you have suffered in the past with cold sores on the skin or lips. This viral infection may be re-activated by some injectable treatments. You are advised to take an anti-viral medication such as Zovirax on the day of your treatment and for several days after your treatment or L-lysine (available from the chemist or health store) for 2-3 days prior to and a week after your treatment.
- If you have also been treated with a skin Booster You will feel raised 'bumps' and raised 'threads' on the skin in some areas that have been treated. It is advisable to massage the areas gently with a greasy moisturiser or arnica cream 2-3 x daily for 5-7 days. This will ensure that the booster is spread across the deep layer of the skin to create a sheet of product which will attract water to it and hydrate the deep layer of the skin.
- DO NOT MASSAGE ANY AREA TREATED WITH PLATELET RICH PLASMA (PRP) or PLATELET RICH PLASMA (PRF) IF YOU HAVE ALSO HAD MUSCLE RELAXANTS INJECTED IN THE SAME AREA. If you feel that further treatment of an area is required, please contact your cosmetic practitioner for a follow-up appointment or advice as to when a further treatment can be carried out.
- If you suffer any allergic responses, prolonged generalised symptoms, Itching, delayed wound healing, signs of infection at the site of blood draw, nerve damage, pain, numbness Please contact us. If any lumps/unevenness persists after 14 days, Please do not hesitate to contact our friendly staff for more information

Profile Clinics 0485 960 001